

Considering OCALIVA[®] (obeticholic acid)

MY HEALTHCARE TEAM DISCUSSION GUIDE

Highlight the questions that apply to you and bring them to your next appointment.

This guide is not intended to replace the advice of your healthcare team. All decisions regarding primary biliary cholangitis (PBC) treatment should be made by the prescriber using his/her clinical judgment.

WHAT IS OCALIVA?

What is OCALIVA and who can take it? Who should not take it?

Can I take OCALIVA if I am taking other medicines for my PBC?
What about taking it with medicines for other health conditions?

Could OCALIVA work for me if I cannot tolerate ursodeoxycholic acid (UDCA)?

What are we trying to achieve by adding OCALIVA to my treatment plan?
Can you help me set short- and long-term goals?

HOW OCALIVA WORKS

Will I be able to tell a difference in how I feel day to day?
How do I know that OCALIVA is working, and how soon will I know?

How is OCALIVA administered? What should I do if I miss a dose?

How does OCALIVA work differently from UDCA?

What dose of OCALIVA is right for me? Will it ever need to be adjusted?

What side effects have patients experienced on OCALIVA?

NOTES

What is OCALIVA?

OCALIVA is a prescription medicine used to treat primary biliary cholangitis (PBC) in combination with ursodeoxycholic acid (UDCA) in adults who have not responded well enough to UDCA, or alone for adults who cannot tolerate UDCA. It is not known if taking OCALIVA will improve your chance of survival or improve your symptoms of PBC. There are ongoing studies to find out how OCALIVA works over a longer period of time.

IMPORTANT SAFETY INFORMATION

What is the most important information I should know about OCALIVA?

OCALIVA may cause serious side effects including:

Worsening of liver problems, liver failure, in some cases leading to death, have happened in people with PBC with advanced liver cirrhosis when OCALIVA was taken more often than recommended.

If you have primary biliary cholangitis (PBC) with advanced cirrhosis, you may need a lower dose of OCALIVA.

Before you start OCALIVA, and during your treatment with OCALIVA, your healthcare provider will do tests to check your liver. These tests will help your healthcare provider decide how much OCALIVA you should take and how often you should take it. If you have worsening liver problems, your dose of OCALIVA may be changed, stopped for a period of time, or stopped completely by your healthcare provider.

Please see additional Important Safety Information on pages 3 and 4 and [Medication Guide](#) and full [Prescribing Information](#) for OCALIVA 5 mg and 10 mg tablets or visit ocaliva.com. Rx only.

HOW TO GET ACCESS TO OCALIVA

How do I receive OCALIVA? How do I set up my initial prescription?
What is a specialty pharmacy? How does it work? How is it different from a regular pharmacy?
How can I find out if my insurance will cover my OCALIVA prescription? How much will OCALIVA cost me?
Who do I call if I need help working with my insurance company?
How can I find out more about financial assistance programs that may cover the cost of OCALIVA?

NOTES

Horizontal lines for taking notes.

RESOURCES AND INFORMATION

How can I find support groups in my area?
How can I learn more about PBC?
How often should I be checking in with my healthcare team?
Should I be seeking advice from other healthcare specialists to manage different aspects of my PBC?

What is OCALIVA?

OCALIVA is a prescription medicine used to treat primary biliary cholangitis (PBC) in combination with ursodeoxycholic acid (UDCA) in adults who have not responded well enough to UDCA, or alone for adults who cannot tolerate UDCA. It is not known if taking OCALIVA will improve your chance of survival or improve your symptoms of PBC. There are ongoing studies to find out how OCALIVA works over a longer period of time.

IMPORTANT SAFETY INFORMATION

What is the most important information I should know about OCALIVA?

OCALIVA may cause serious side effects including:

Worsening of liver problems, liver failure, in some cases leading to death, have happened in people with PBC with advanced liver cirrhosis when OCALIVA was taken more often than recommended.

Please see additional Important Safety Information on pages 3 and 4 and [Medication Guide](#) and full [Prescribing Information](#) for OCALIVA 5 mg and 10 mg tablets or visit ocaliva.com. Rx only.



IMPORTANT SAFETY INFORMATION (CONT'D)

What is the most important information I should know about OCALIVA?

If you have primary biliary cholangitis (PBC) with advanced cirrhosis, you may need a lower dose of OCALIVA. Before you start OCALIVA, and during your treatment with OCALIVA, your healthcare provider will do tests to check your liver. These tests will help your healthcare provider decide how much OCALIVA you should take and how often you should take it. If you have worsening liver problems, your dose of OCALIVA may be changed, stopped for a period of time, or stopped completely by your healthcare provider.

Tell your healthcare provider right away if you have any of the following symptoms of worsening liver problems during treatment with OCALIVA:

- Swelling of your stomach area from a build-up of fluid; yellowing of your skin or the whites of your eyes; black, tarry, or bloody stools; coughing up or vomiting blood, or your vomit looks like “coffee grounds”; or mental changes (such as confusion, sleepier than usual or harder to wake up, slurred speech, mood swings, or changes in personality)

Tell your healthcare provider right away if you have any of the following symptoms during treatment with OCALIVA **and they are severe or do not go away:**

- Stomach-area pain, nausea, vomiting, or diarrhea; loss of appetite or weight loss; new or worsening fatigue, weakness, fever, or chills; light-headedness; less frequent urination

Who should not take OCALIVA?

Do not take OCALIVA if you have or had a complete blockage in the bile ducts in your liver or gallbladder.

What are the possible side effects of OCALIVA?

OCALIVA may cause serious side effects including:

- See “**What is the most important information I should know about OCALIVA?**”
- **Severe Itching.** Itching (pruritus) is a common side effect and can sometimes become severe (intense itching or itching all over your body). Severe itching can cause discomfort, problems sleeping, and problems doing daily activities, and usually needs to be treated. Tell your healthcare provider if you get severe itching or if your itching gets worse.
- **Decreases in Good Cholesterol.** Decreases in HDL-C (“good cholesterol”) have been observed in patients taking OCALIVA. Your healthcare provider will check your cholesterol levels during treatment to see if you should continue taking OCALIVA.

The most common side effects of OCALIVA include: pruritus (itching of the skin), tiredness, stomach pain and discomfort, rash, joint pain, mouth and throat pain, dizziness, constipation, swelling in your hands, ankles or feet, fast or irregular heartbeat, fever, changes in how your thyroid gland works, and eczema (skin dryness, irritation, redness, crusting, or drainage).

These are not all the possible side effects associated with OCALIVA. Call your healthcare provider for medical advice about side effects.

What should I tell my healthcare provider before taking OCALIVA?

Before taking OCALIVA, tell your healthcare provider about all of your medical conditions, including if you:

- **are pregnant or plan to become pregnant.** It is not known if OCALIVA will harm your unborn baby.
- **are breastfeeding or plan to breastfeed.** It is not known if OCALIVA passes into your breastmilk. Talk with your healthcare provider about the best way to feed your baby if you take OCALIVA.

Tell your healthcare provider about all the medicines you take, including prescription and over-the-counter medicines, vitamins, and herbal supplements. OCALIVA can affect the way certain medicines work. Certain other medicines may affect the way OCALIVA works.

Please see [Medication Guide](#) and full [Prescribing Information](#) for OCALIVA 5 mg and 10 mg tablets or visit ocaliva.com.

Available by prescription only.

To report negative side effects of OCALIVA, please contact Intercept Pharmaceuticals, Inc. at 1-844-782-ICPT or you may report to FDA at 1-800-FDA-1088 or www.fda.gov/medwatch.